

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/00, 7/00, C12P 21/00	A1	(11) International Publication Number: WO 90/09441 (43) International Publication Date: 23 August 1990 (23.08.90)
---	----	--

(21) International Application Number: PCT/US90/00357
(22) International Filing Date: 22 January 1990 (22.01.90)
(30) Priority data:
304,619 1 February 1989 (01.02.89) US

(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; Fruit Street, Boston, MA 02114 (US).

(72) Inventors: GELLER, Alfred, I. ; 28 Holbrook Street, Jamaica Plains, MA 02130 (US). BREAKFIELD, Xandra, O. ; 127 Homer Street, Newton, MA 02159 (US).

(74) Agents: GOLDSTEIN, Jorge, A. et al.; Saidman, Sterne, Kessler & Goldstein, 1225 Connecticut Avenue, N.W., Suite 300, Washington, DC 20036 (US).


(81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: HERPES SIMPLEX VIRUS TYPE I EXPRESSION VECTOR



(57) Abstract

A recombinant HSV-1 vector capable of infecting and being propagated in a non-mitotic cell. The vector can be used to treat neurological diseases, and to produce animal and *in vitro* models of such diseases.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LJ	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

TITLE OF THE INVENTION

HERPES SIMPLEX VIRUS TYPE I EXPRESSION VECTOR

FIELD OF THE INVENTION:

5 The present invention relates to a defective herpes simplex type I vector capable of expressing a gene product in eukaryotic cells. The invention further pertains to the use of this vector in introducing and expressing gene sequences in post-mitotic cells such as neurons and in other cells of the nervous system. This invention was made with Federal funds, the Government has certain rights in this invention.

10 BACKGROUND OF THE INVENTION

15 The capacities to introduce a particular foreign or native gene sequence into a mammalian cell and to control the expression of that gene are of substantial value in the fields of medical and biological research. Such capacities provide a means for studying gene regulation, for defining the molecular basis of a disease, and for designing a therapeutic basis for the treatment of disease.

20 Gene transfer techniques can be used for two general purposes. First, they can be used to evaluate the regulation and function of a cloned gene following its modification and introduction into different cell types by studying, for example, the definition of the regulatory elements that control levels of gene expression, alternate modes of RNA splicing, post-translational processing of peptides, sorting

-2-

of proteins to their appropriate cellular locations, and biological activities of proteins. Second, gene transfer techniques can be used to modify cells, such as those of the nervous system, in culture and in vivo. Such studies may, for example, involve analysis of cell lineage, alternation of phenotypic properties, and ablation of specific cell populations, as well as creation and correction of hereditary disease states. Gene transfer techniques can thus be used as a tool in understanding molecular aspects of the development, function, and survival of cells.

The introduction of a particular foreign or native gene into a mammalian host cell is facilitated by first introducing the gene sequence into a suitable nucleic acid vector. A variety of methods have been developed which are capable of permitting the introduction of such a recombinant vector into a desired host cell.

For example, such a recombinant vector can be introduced into the host cell by DNA-mediated transformation (Choo, K.H. et al., Gene 46:277-286 (1986); Perez, C.F. et al., Radiat. Res. 104:200-213 (1985); Horst, J. et al., Hoppey-Seyler's Z. Physiol. Chem. 363:445-448 (1982); Hirschhorn, R.R. et al., Fed. Proc. 41: Abstract 6525 (1982); Graham, F.L. et al., In: The Wistar Symposium Series, Volume 1 Introduction of Macromolecules into Viable Mammalian Cells, Alan R. Liss, Inc., New York, N.Y. page 3-26 (1980); Upcroft, P., Anal. Biochem. 162:1-4 (1987)). The vector can also be introduced into a mammalian cell by protoplast fusion (Yoakum, G.H., Biotechniques 2:24-26, 28-30 (1984)), or by micro-injection (Spandidos, D.A. et al., Eur. J. Cell. Biol. 37:234-239 (1985); Folger, K.R. et al., Molec. Cell. Biol. 2:1372-1387 (1982); Gordon, J.W. et al., Proc. Natl Acad. Sci. USA 77:7380-7384 (1980)). Unfortunately, the above-described techniques are relatively inefficient and unsuitable for use in situations which require that the recombinant molecule be

-3-

introduced into all or most of the cells present in culture or in an animal.

5 Techniques of transgenic genetics have been used to achieve the efficient introduction of a cloned gene sequence into all or most of the cells of an animal. In such an approach, a recombinant plasmid is introduced into the pronuclei of a fertilized egg and permitted to develop into a transgenic animal. Such an animal will usually contain the introduced gene in all of the cells of its body including its germ line. Transgenic genetics is, however, a technically difficult and exacting procedure.

10 Viral vectors have been employed in order to increase the efficiency of introducing a recombinant vector into suitably sensitive host cells. Viruses which have been employed as vectors for the transduction and expression of exogenous genes in mammalian cells include SV40 virus (Chung, M.H. et al., Korean J. Microbiol. 25:165-172 (1987); Innis, J.W. et al., Molec. Cell. Biol. 3:2203-2210 (1983); Okayama, H. et al., Molec. Cell. Biol. 5:1136-1142 (1985)), bovine papilloma virus (Meneguzzi, G. et al., Embo. J. 3:365-372 (1984); Dimaio, D. et al., Proc. Natl. Acad. Sci. USA 79:4030-4034 (1982); Lusk, M. et al., Cell 36:391-402 (1984); Giri, I. et al., Virology 127:385-396 (1983); Lusk, M. et al., Molec. Cell. Biol. 3:1108-1122 (1983)), etc.

25 Retroviruses which have been employed as vectors for the transduction and expression of exogenous genes in mammalian cells include the Moloney murine sarcoma virus (Perkins, A.S. et al., Molec. Cell. Biol. 3:1123-1132 (1983); Lee, W.H. et al., J. Virol. 44:401-412 (1982); Curran, T. et al., J. Virol. 44:674-682 (1982); Gazit, A. et al., J. Virol. 60:19-28 (1986)), etc. In contrast to methods which involve DNA transformation or transfection, the use of viral vectors can result in the rapid introduction of the recombinant molecule into a wide variety of host cells.

-4-

Efforts to introduce recombinant molecules into post-mitotic neurons and other neural cells have, however, been hampered by the inability of such cells to be infected by the above-described viral or retroviral vectors. Thus, the study of gene expression in neuronal cells, and the identification of therapies for treating neuronal disease have been hampered by the lack of suitable methods to accomplish gene transfer into neural cells. A need therefore exists for efficient viral vectors capable of mediating gene transfer into such cells.

SUMMARY OF THE INVENTION

The invention concerns a recombinant HSV-1 vector capable of infecting and persisting in a non-mitotic cell. The vector can be used to treat neurological diseases, and to produce animal and in vitro models of such diseases. In detail, the invention provides a recombinant HSV-1 vector capable of infecting a neuronal cell, wherein the vector contains:

- (1) an HSV-1 packaging site-containing sequence;
- (2) an HSV-1 origin of DNA replication-containing sequence; and
- (3) a heterologous reporter or selectable marker gene sequence sufficient to permit the recognition or selection of the vector.

The invention also concerns the above recombinant vector which additionally contains one or more of the following sequences:

- (4) a promoter sequence sufficient to direct transcription of a distally located sequence in a neuronal cell;
- (5) a sequence sufficient to direct translation of a distally located sequence in a neuronal cell;

-5-

- 5
- (6) a sequence sufficient to permit a neuronal cell to process an expressed protein;
 - (7) a sequence sufficient to permit the propagation of the vector in a bacteria; or
 - (8) an inserted gene sequence wherein the inserted gene sequence is inserted into the vector such that it is operably linked to the sequences (4), (5) and (6).

10 The invention further concerns the above-described recombinant vectors which are capable of persisting in a non-mitotic cell.

The invention further concerns the above-described recombinant vector which is pHSVlac.

15 The invention further concerns the above-described recombinant vector wherein the inserted gene sequence (8) is a gene sequence associated with a disease selected from the group consisting of: lysosomal storage disease, the Lesch-Nyhan syndrome, amyloid polyneuropathy, Alzheimer amyloid, and Duchenne's muscular dystrophy.

20 The invention further concerns a method for expressing a gene sequence in a non-mitotic cell which comprises:

- (a) operably linking the gene sequence to a promoter sequence of a recombinant vector capable of persisting in the non-mitotic cell;
- 25 (b) introducing the vector into the cell; and
- (c) permitting the vector to express the gene in the cell.

The invention further concerns a method for treating a neurological deficiency state disease which comprises:

- 30
- (a) operably linking a gene sequence whose gene product complements the deficiency of the deficiency state disease to a promoter sequence of a recombinant vector capable of propagation in a neuronal cell;
 - (b) introducing the vector into the cell; and

-6-

(c) permitting the vector to express the gene in the cell, wherein the expression complements the deficiency.

The invention further concerns a method for identifying an agent capable of treating a neurological unbalanced state disease which comprises:

(a) operably linking a gene sequence whose expression is sufficient to induce the unbalanced state disease in an animal to a promoter sequence of a recombinant vector capable of propagation in a neuronal cell;

(b) introducing the vector into the cell; and

(c) permitting the vector to express the gene in the cell, wherein the expression induces the unbalanced state disease;

(d) providing to the cell an agent suspected of being able to treat the unbalanced state disease in the cell; and

(e) identifying any agent capable of treating the unbalanced state disease in the cell.

BRIEF DESCRIPTION OF THE FIGURE

Figure 1 shows the structure of pHSVlac. The clear region contains the HSV-1 a segment nucleotides 127 to 1132, the packaging site (Davison, A.J. et al., J. Gen. Virol. 55:315 (1981)). The cross hatched region symbolizes the HSV-1 c region, nucleotides 47 to 1066 (McGeoch, D.J. et al., Nucleic Acids Res. 14:1727 (1986)). pHSVlac was constructed from pCH110 (Hall, C.V. et al., J. Molec. App. Genet. 2:101 (1983)).

DESCRIPTION OF THE PREFERRED EMBODIMENTSI. Methods for Introducing Gene Sequences into Neuronal and Other Post-Mitotic Cells

5 The present invention concerns a means for introducing gene sequences into cells of the nervous system such as neurons, neuroglial cells, etc. Such cells are collectively described herein as "neural or neuronal" cells. Neural cells are described, for example, by Barr, M.L., The Human Nervous System An Anatomic Viewpoint, 3rd. Ed., Harper & Row, NY
10 (1979), which reference is herein incorporated by reference).

To study the functions of cloned neuronal genes, methods are required to deliver genes into the cells of the nervous system. Of the four approaches used to introduce genes into cells (the frog oocyte micro-injection system (Noda, M. et al., Nature 302:818 (1983)), transgenic mice (Palmiter, R.D. et al., Science 222:809 (1983)), transfection of DNA directly into cells (Graham, F.L. et al., Virology 52:456 (1973)), and retrovirus vectors (Mann R. et al., Cell 33:153 (1983)) none
15 can deliver a gene directly into non-mitotic cells.

20 Although delivery of recombinant molecules into the cells of the nervous system prior to birth can be accomplished using transformation or micro-injection, delivery into such cells after birth presents complications because of the relative inaccessibility of the cells, their diversity, and the post mitotic state of mature neurons. Thus, such methods are
25 relatively unsuitable for introducing gene sequences into neuronal cells.

In view of the above-stated deficiencies of existing techniques for introducing gene sequences into a neural cell,
30 it would be highly desirable to develop suitable viral vectors which could be employed to deliver such sequences to neural cells. The development of such a gene sequence delivery

system is one aspect of the present invention. Methods for introducing gene sequences into neuronal cells are reviewed by Breakefield, X.O. *et al.*, Molec. Neurobiol. 1:339-371 (1987), which reference is herein incorporated by reference in its entirety).

A viral vector, as that term is used herein, is a nucleic acid molecule (preferably of DNA) in which a gene sequence (which is to be transferred) is fused to a subset of viral sequences. The viral sequences and the total genome size is selected such that the vector is capable of being encapsulated in a virus particle and thus be capable of binding to, and introducing its gene sequences into a virus-sensitive host cell. The infective properties of such a virion are, thus, the same as those containing the wild type viral genome.

The term "gene sequence," as used herein, is intended to refer to a nucleic acid molecule (preferably DNA). Such gene sequences may be derived from a variety of sources including DNA, cDNA, synthetic DNA, RNA, or combinations thereof. Such gene sequences may comprise genomic DNA which may or may not include naturally occurring introns. Moreover, such genomic DNA may be obtained in association with promoter regions or poly A sequences. The gene sequences of the present invention are preferably cDNA. Genomic DNA or cDNA may be obtained in any of several ways. Genomic DNA can be extracted and purified from suitable cells by means well-known in the art. Alternatively, mRNA can be isolated from a cell and used to produce cDNA by reverse transcription, or other means.

In order to produce the viral vectors of the present invention, several factors are preferably considered. First, since the vector will be delivered only into those cells that can be infected (either abortively or productively) by the wild type virus, it is necessary to produce the viral vector from a wild type virus which is capable of infecting neuronal

-9-

cells. One must also consider the ability of the viral vector to infect cells in tissue culture or in vivo.

5 Second, since it may be desirable to maintain the vector and to express the cloned gene sequence for an extended period of time, the virus employed should be capable of persisting in a cell for an extended period of time without causing cell death.

10 Third, since viral vectors can be produced from viruses which are maintained in either the nucleus or the cytoplasm of a neuronal cell, one must consider whether one desires the viral gene fusion to be maintained in the nucleus or in a cytoplasm of the recipient cell. If, for example, the viral vector exists as a double stranded DNA molecule in the nucleus of a cell, than its transcription may retain appropriate
15 responses to cellular trans-acting regulatory factors. In contrast, if a viral vector is maintained in the cytoplasm, it may not encounter such trans-acting regulatory factors, and therefore, not be subjected to regulation.

20 Fourth, since the size of the virus genome determines an upper limit on the size of the vector and the gene sequences that can be inserted in it, it is necessary to produce the viral vector from a virus whose size is sufficiently large that it may be adapted to contain an inserted gene sequence of desired size.

25 In selecting a viral vector system capable of delivering gene sequences to neuronal cells, it has been found to be most desirable to employ a DNA virus whose site of replication is in the nucleus of infected cells. It has further been found desirable for the virus to be capable of latent infection
30 (i.e., persisting in an infected cell without causing cell death). Preferably, the viral vector should also be of substantial size in order that viral derivatives can be formed which are capable of containing substantial amounts of introduced gene sequences. The attributes of such a virus

-10-

have been found to be present in Herpes Simplex Virus-1 ("HSV-1") (Breakefield, X.O. et al., Mol. Neurobiol. 1:339 (1987); Geller, A.I. et al. Science:241:1667- 1669 (1988), which reference is herein incorporated by reference in its entirety).

HERPES SIMPLEX VIRUS - 1

HSV-1 is a double-stranded DNA virus which is replicated and transcribed in the nucleus of the cell. In virions, HSV-1 vectors are composed of head to tail repeats (Stow, N.D. et al., Eukaryotic Viral Vectors, Y. Gluzman, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982), pp. 199-204; Spaete, R.R. et al., Cell 30:285 (1982)). The genes of HSV-1 are transcribed by the cellular RNA polymerase II. HSV-1 contains approximately 100 genes and is approximately 150 kilobases in size. Five immediate early ("IE") genes encode the major regulatory proteins of the virus. Immediate early genes induce expression of the early ("E") genes that are responsible for DNA replication. The late ("L") genes are induced after DNA replication and encode the structural components and enzymes required for assembly of virus particles. When the late genes are induced, transcription of the immediate early genes is reduced.

HSV-1 possesses both a lytic and a latent cycle. During the lytic cycle of HSV-1, expression of the three classes of genes occurs in a complex sequential cascade. In contrast, in its latent cycle, HSV-1 gene expression is limited to at most the IE genes and a latency associated transcript (Wagner, E.K. et al., Science 235:1056 (1987); Deatly, A.M. et al., Proc. Natl. Acad. Sci. USA 84:3204 (1987); Wagner, E.K. et al., J. Virol. 62:1194 (1988); Green, M.T. et al., Infect. Immunol. 34:987 (1981); Persson, R.H. et al., J. Virol. 54:414 (1985); Mosca, J.D. et al., J. Virol. 56:867 (1985)). DNA replication

-11-

does not occur and no progeny virus are produced (Stevens, J.G., Current Topics in Microbiology and Immunology 70:31 (1975)).

5 Thus, in both lytic and latent states, extensive gene regulation occurs which closely mimics the regulation of cellular genes. Electrophysiological properties are unaltered in latently infected neurons (Fukuda, J. et al., Brain Res. 262:79 (1983)).

10 HSV-1 has a wide host range, and infects many cell types in mammals and birds (including chickens, rats, mice, monkeys, and humans) (Spear, P.G. et al., DNA Tumor Viruses, J. Tooze, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1981), pp. 615-746). HSV-1 can lytically infect a wide variety of cells including neurons, fibroblasts, and
15 macrophages. In addition, HSV-1 infects post-mitotic neurons in adult animals and can be maintained indefinitely in a latent state (Stevens, J.G., Current Topics in Microbiology and Immunology 70:31 (1975)).

20 Post-mitotic neurons harbor HSV-1 in the virus' latent state (Stevens, J.G., Curr. Top. Microbiol. Immunol. 70:31-50 (1975)). Once HSV-1 attains latency, it can be retained for the life of the neuron. As mentioned above, latent HSV-1 is capable of expressing genes; expression of genes encoded by HSV-1 has been detected by immunochemistry in latently
25 infected neurons. Furthermore, HSV-1 is transported both anterogradely and retrogradely in neurons. This property is especially advantageous in that it suggests that HSV-1 vectors will be capable of reaching cells of interest some distance away from the injection site.

30 Among the immortal neuronal cell lines that can be infected by HSV-1 include the mouse neuroblastoma cell lines NS20Y and N1E-115, and the rat pheochromocytoma cell line PC12. HSV-1 virions can be detected in neurons in vivo by electron microscopy (Cook, M.L. et al., Infect. Immunity.

-12-

9:946-951 (1974)). Two lines of evidence suggest that HSV-1 can infect most, if not all, kinds of neurons in the central nervous system. First, following inoculation of HSV-1 in the periphery, a burst of virus production ascends the neuroaxis, initially in the sensory or motor neurons innervating the site of inoculation, then in the spinal cord, brain stem, cerebellum, and cerebral cortex (Koprowski, H., In: Persistent Viruses (Stevens, F.G., ed.), pp. 691-699, Academic Press, NY (1978)). Second, attempts to mimic HSV-1 latency in tissue culture with different preparations of neurons have required high temperature, DNA synthesis inhibitors, and antisera directed against HSV-1 virions to prevent a lytic infection for spreading to all the neurons (Wigdahl, B., et al., Proc. Natl. Acad. Sci. USA 81:6217-6201 (1984)).

THE HSV-1 VECTORS OF THE PRESENT INVENTION

The preferred vectors of the present invention are produced from HSV-1. The HSV-1 vectors of the present invention will preferably contain:

(1) a sequence which contains an HSV-1 packaging site (preferably the HSV-1 α segment located at approximately nucleotides 127-1132 of the α sequence of HSV-1 virus, or its equivalent) (Davison, A.J. et al., J. Gen. Virol. 55:315 (1981) such that the vector can be packaged into a particle which is capable of adsorbing to a cell;

(2) a sequence which contains an HSV-1 origin of DNA replication (preferably the HSV-1 γ region, containing the HSV-1 ori_S region, located at approximately nucleotides 47-1066 of HSV-1 virus, or its equivalent) (McGeoch, D.J. et al., Nucleic Acids Res. 14:1727 (1986)); and

(3) a reporter or selectable marker gene sequence sufficient to permit the recognition or selection of the

-13-

vector in a host cell. A reporter gene sequence, as used herein, is any gene sequence which, when expressed, results in the production of a protein whose presence or activity can be easily monitored. As used herein, a reporter or selectable marker gene sequence is said to be "heterologous" if it is not naturally present in a wild type HSV-1 genome. Examples of suitable reporter genes include the gene for galactokinase, beta-galactosidase, chloramphenicol acetyltransferase, beta-lactamase, etc. Alternatively, the reporter gene sequence may be any gene sequence whose expression produces a gene product which affects neuronal physiology (including components of second messenger systems or neurotransmitter metabolism, or ion channels).

A selectable marker gene sequence is any gene sequence capable of expressing a protein whose presence permits one to selectively propagate a cell which contains it. Examples of selectable marker gene sequences include gene sequences capable of conferring host resistance to antibiotics (such as ampicillin, tetracycline, kanamycin, streptomycin, etc.), of conferring resistance to amino acid analogues, of permitting the growth of the bacteria on additional carbon sources or under otherwise impermissive culturing conditions. A gene sequence may be both a reporter gene sequence and a selectable marker gene sequence; The most preferred reporter gene of the present invention is the lacZ gene which encodes the beta-galactosidase activity of E. coli. The most preferred selectable marker sequence is the beta-lactamase gene of pBR322;

The most preferred reporter or selectable marker gene sequence sufficient to permit the recognition or selection of the vector in a neuronal cell. Preferably, the selected reporter gene sequence will encode an enzyme or other protein which is normally absent from mammalian cells, and whose presence can, therefore, definitively establish the presence

-14-

of the vector in such a cell. The most preferred such sequence being the E. coli lacZ gene (Sanes, J.R. *et al.*, EMBO J. 5:3133 (1986); Price, J. *et al.*, Proc. Natl. Acad. Sci. USA 84:156 (1987)).

5 An even more preferred vector of the present invention will, in addition to the above-enumerated sequences (1), (2) and (3), contain one or more of the following sequences:

10 (4) a sequence, such as a promoter sequence, sufficient to direct the transcription of a distally located sequence (i.e. a sequence linked to the 5' end of the promoter sequence) in a neuronal cell. The most preferred such sequence being the HSV-1 IE 4/5 promoter sequence (McGeoch, D.J. *et al.*, Nucleic Acids Res. 14:1727 (1986)).

15 (5) a sequence, such as an ATG or other initiation codon, sufficient to direct the translation of the distally located sequence in a neuronal cell. The most preferred such sequence being the sequence which follows the HSV-1 IE 4/5 promoter sequence;

20 (6) a sequence, such as a polyadenylation sequence, a localization sequence, a signal sequence, or a termination sequence, sufficient to permit a neuronal cell to efficiently and effectively process the expressed protein. The most preferred polyadenylation sequence being the SV40 early region polyadenylation site (Hall, C.V. *et al.*, J. Molec. App. Genet. 2:101 (1983));

25 (7) a sequence sufficient to permit the propagation of the viral vector in a bacteria, such as E. coli. Such a sequence may comprise only an origin of replication, or may comprise an entire plasmid capable of replicating in the
30 bacteria. For example, in a preferred embodiment, the bacteria is E. coli, and the sequence can be either an E. coli ori sequence or (even more preferably) a plasmid such as Col E1, pBR322, etc. The most preferred such sequence contains

-15-

the Col E1 origin of replication of pBR322. The entire pBR322 plasmid may be employed;

An even more preferred vector of the present invention will, in addition to one or more of the above-enumerated sequences, contain:

5 (8) one or more inserted sequences whose introduction, transcription, or expression in a neuronal cell is desired. Such sequences are inserted into the vector in order to operably link the sequence with the promoter sequence (4), if
10 transcription is desired, or additionally with the initiation and processing sequences (5 and 6), if translation and processing are desired. Alternatively, the inserted sequence may be placed at any position in the vector. The term
15 "operable linkage" is intended to describe a linkage between a gene sequence and a promoter or other regulatory or processing sequence such that the transcription of the gene sequence is directed by an operably linked promoter sequence, the translation of the gene sequence is directed by an operably
20 linked translational regulatory sequence, and the post-translational processing of the gene sequence is directed by an operably linked processing sequence.

It is, of course possible to employ, as the inserted gene sequence a gene sequence which already possesses a promoter, initiation sequence, or processing sequence. In such a case,
25 it would be unnecessary to employ a viral vector which additionally contained these sequences.

As will be appreciated by one of ordinary skill, the nucleotide sequence(s) of the inserted gene sequence or sequences may be of any nucleotide sequence.

30 The viral vectors of the present invention are preferably from 5 to 15 kb in size, and encapsulated in the HSV-1 viral particle as head to tail repeats of varying length up to 150 kb (the size of wild type HSV-1). HSV-1 contains three origins of DNA replication (ori), or one ori every 50 kb,

-16-

while the above-described vector contains one origin of DNA replication every 5 kb to 15 kb (Stow, N.D. et al., Eukaryotic Viral Vectors, Y. Gluzman, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982), pp. 199-204; Spaete, R.R. et al., Cell 30:285 (1982)).

Viral stock containing any of the viral vectors of the present invention can be preferably obtained by transfecting fibroblasts with a mixture of vector DNA and helper HSV-1 DNA (Stow, N.D. et al., Eukaryotic Viral Vectors, Y. Gluzman, Ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1982), pp. 199-204; Spaete, R.R. et al., Cell 30:285 (1982)). In such an infection, the vectors are maintained and propagated because their possession of an increased number of origins of replication provide them with a growth advantage over the helper virus (Frenkel, N., et al., Ann. N.Y. Acad. Sci. 354:347-370 (1980)). Alternatively, the vector can be propagated according to the manner of Geller, A.I. (Nucl. Acid Res. 16:5690 (1988)).

Although it is possible to propagate the HSV-1 viral vector using a wild type HSV-1 virus (as a helper virus), infection of neuronal cells with wt HSV-1 invariably causes some cell death. Since such cell death is especially undesirable for many gene expression studies, it is especially preferred to propagate the HSV-1 vector using an HSV-1 helper virus which has a mutation, most preferably a temperature sensitive mutation, which prevents the helper virus from entering the lytic cycle (thereby preventing cell damage) when the neuronal cell infection is carried out at a viral growth restrictive temperature of 37 or 39 °C. Such a helper virus can be grown at the viral growth permissive temperature of 31°C.

Intracerebral injection (Watson, K. et al., J. Gen. Virol. 49:149 (1980)) and infection of mouse neuroblastoma cells (Gerdes, J.G. et al., Virology 94:430 (1979)) with HSV-1

-17-

temperature sensitive (ts) mutants allows persistence of the virus without cell death.

Structural and genetic characteristics of the HSV-1 vectors of the present invention can be most clearly illustrated by a description of the most preferred HSV-1 vector: pHSVlac. This defective HSV-1 viral vector contains 8.1 kb of double stranded DNA. The vector is depicted in Figure 1. pHSVlac contains the following gene sequences:

- (1) an HSV-1 packaging site comprising approximately nucleotides 127-1132 of HSV-1 a region;
- (2) an HSV-1 origin of DNA replication comprising the HSV-1 c region located at approximately nucleotides 47-1066;
- (3) the beta-lactamase gene sequence of pBR322; and
- (4) the HSV-1 IE 4/5 promoter sequence;
- (5) the intervening sequence (intron) following the HSV-1 IE 4/5 promoter sequence;
- (6) the SV40 early region polyadenylation sequence;
- (7) the Col E1 origin of replication region of pBR322; and
- (8) the E. coli lacZ gene (as an inserted gene sequence whose expression in a neuronal cell is readily detectable).

pHSVlac was deposited with the American Type Culture Collection on January 31, 1989, and given the accession number ATCC 40544.

pHSVlac DNA is preferably packaged into HSV-1 virus particles using HSV-1 strain 17 ts K (Davison, M.J. et al., J. Gen. Virol., 65:859 (1984); Brown, S.M. et al., Proc. Natl. Acad. Sci. (U.S.A.) 76:2364-2368 (1979); Watson, K., J. Gen. Virol. 49:149-160 (1980); Dargan, D. et al., J. Gen. Virol. 64:1311-1326 (1983); Matz, B. et al., J. Gen. Virol. 64:2261-

-18-

2270 (1983); Dargan, D. et al., J. Gen. Virol. 65:477-492 (1984) as helper virus (hereinafter referred to as "ts K"). HSV-1 strain 17 ts K has a mutation in the IE 3 gene. The virus has an immediate early phenotype, and is not permissive for DNA replication at 37-39° C. It can, however, replicate when cultured at 31° C.

pHSVlac contains three Eco RI sites, one at each end of the pBR322 segment and a third in the lacZ gene 133 bp from the 3' end of the gene. Most of the transcription unit of pHSVlac is present in a 4.3 kb fragment. The pBR322 sequences of pHSVlac are present in a 2.3 kb fragment. The 3' end of the lacZ gene, the SV-40 early region polyadenylation site, and the HSV-1 a sequence are contained in a 1.5 kb fragment of pHSVlac.

APPLICATIONS OF GENE TRANSFER TECHNOLOGY TO NEUROSCIENCE AND NEUROCHEMISTRY

Gene transfer technology has several applications to neuroscience and neurochemistry. The most immediate applications are, perhaps, in elucidating the process of neural peptides and the functional domains of proteins. Cloned cDNA or genomic sequences for neural proteins can be introduced into different cell types in culture, or in vivo, in order to study cell type-specific differences in processing and cellular fate. By placing the coding sequences under the control of a strong promoter, a substantial amount of the protein can be made, thus avoiding difficulties in characterizing trace amounts. Furthermore, the specific residues involved in protein processing, intracellular sorting, or biological activity can be determined by mutational change in discrete residues of the coding sequence.

Gene transfer technology can also be applied to provide a method to control expression of a protein and to assess its

capacity to modulate cellular events. Some functions of neural proteins, such as their role in differentiation, may be studied in tissue culture, whereas others will require re-introduction into the nervous system at different times in development or aging in order to monitor changes in receptor density, cell number, fiber growth, electrical activity, and other relevant properties.

Gene transfer provides a means to study the DNA sequences and cellular factors which regulate expression of neural specific genes. One approach to such a study would be to fuse the regulatory elements to be studied to a particular reporter gene and subsequently assaying for the expression of the reporter gene.

The regulation of gene expression in neuronal cells has been found to have a role in maintaining homeostasis and is believed to have a role in mediating information retention in response to external and internal signals (Black, I.B., et al., Science 236:1263-1268 (1987)). During development, coordinate regulation of gene expression serves to produce a differentiated phenotype, e.g., as in catecholamine metabolism and myelin biosynthesis. Regulation depends on many factors including chromatin structure, DNA methylation, and trans-acting factors, which respond to phosphorylation, hormones, and other signals. It is a complex process that allows sets of genes to be expressed together or differentially and may involve a combinatorial code of regulatory sequences.

Issues of cellular fate and interactions in the nervous system can also be addressed by gene transfer. For example, genes which encode histological markers can be introduced into embryonic cells to determine lineage relationships during development and to elucidate neuronal pathways. In addition, genes encoding growth factors, oncogenic proteins, toxic peptides, or other physiologically important proteins, can be introduced into specific areas of the nervous system to study

-20-

their effects on cell division, survival, and differentiation. For some studies, gene transfer or gene expression must be restricted to specific cells in the nervous system.

5 Gene transfer also possesses substantial potential use in understanding and providing therapy for disease states. There are a number of inherited neurologic diseases in which defective genes are known and have been cloned. In some cases, the function of these cloned genes is known. In
10 humans, genes for defective enzymes have been identified for (1) lysosomal storage diseases such as those involving β -hexosaminidase (Kornerluk, R.G., et al., J. Biol. Chem. 261:8407-8413 (1986); Myerowitz, R., et al., Proc. Natl. Acad. Sci. USA 82:7830-7834 (1985)) and glucocerebrosidase (Sorge et al., Proc. Natl. Acad. Sci. USA 82:5442-5445 (1985); Tsuji, S., et al., N. Engl. J. Med. 316:570-575 (1987)), (2) for deficiencies in hypoxanthine phosphoribosyl transferase activity (the "Lesch-Nyhan" syndrome; Stout et al., Met. Enzymol. 151:519-530 (1987)), (3) for amyloid
15 polyneuropathies (prealbumin; Sasaki, H., et al., Biochem. Biophys. Res. Commun. 125:636-642 (1984), (4) for Alzheimer amyloid (Tanzi, R.E., et al., Science 235:880-884 (1987); Goldgaber, D., et al., Science 235:877-880 (1986)); (5) for Duchenne's muscular dystrophy (uncharacterized muscle protein; Monaco, A.P., et al., Nature 323:646-650 (1987)); and (6) for
20 retinoblastoma (uncharacterized protein expressed in the retina and other tissues, Lee, W.-H., et al., Science 235:1394-1399 (1987); Friend, S.H., et al., Nature 323:643-646 (1986)).

30 Gene transfer techniques can also be used to study the "shiverer" mutation (myelin basic protein, Roach, A., et al., Cell 42:149-155 (1987); Molineaux, S.M., et al., Proc. Natl. Acad. Sci. USA 83:7542-7546 (1986)) and the "jimpy" mutation (proteolipoprotein, Nave, K.-A., et al., Proc. Natl. Acad.

-21-

Sci. USA 83:9264-9268 (1986); Hudson, L.D., et al., Proc. Natl. Acad. Sci. USA 84:1454-1458 (1987)).

5 The above diseases fall into two classes: deficiency states, usually of enzymes, which are inherited in a recessive manner; and unbalanced states, at least sometimes involving structural or regulatory proteins, which are inherited in a dominant manner.

10 For deficiency state diseases, gene transfer could be used to bring a normal gene into affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For unbalanced state diseases, gene transfer could be used to create the disease state in a model system, which could be used in efforts to counteract the effect of the imbalance. Thus, the methods of
15 the present invention permit the treatment of neurological diseases. As used herein, a deficiency state disease is "treated" by partially or wholly remedying the deficiency which causes the deficiency or which makes it more severe. As used herein, an unbalanced state disease is "treated" by
20 partially or wholly remedying the imbalance which causes the disease or which makes it more severe. The use of site-specific integration of DNA sequences to cause mutations or to correct defects is also possible.

25 In summary, HSV-1 viral vectors, such as pHSVlac, can be used to stably transfect post-mitotic cells such as neurons and stably express β -galactosidase. The E. coli lacZ gene is pHSVlac can be exchanged for other coding sequences. Using this vector, it is now possible to introduce, into neurons, genes which encode products that effect physiology, including
30 components of second messenger systems and neurotransmitter metabolism. The vectors of the present invention may be useful for gene therapy on neuronal diseases or for experimental modification of neuronal physiology.

-22-

Having now generally described this invention, the same will be better understood by reference to certain specific examples which are included herein for purposes of illustration only and are not intended to be limiting of the invention, unless specified.

EXAMPLE 1

METHODS FOR CELL AND VIRAL PROPAGATION

CV1 monkey fibroblasts were grown in Dulbecco's modified minimum essential medium with 10% fetal calf serum. HSV-1 strain 17 ts K was describe by Davison, M.J. et al., J. Gen. Virol., 65:859 (1984)). Cultures of embryonic mouse spinal cord were obtained according to the procedure of Huettnner, J.E. et al. (J. Neurosci. 6:3044-3066 (1986)). Cultures of striatum and total neocortex (Freese A. et al., Proc. Soc. Neurosci. 14:Abstract 169.10 (1988)), cerebellum, thalamus, and cortical areas (Huettnner, J.E. et al., J. Neurosci. 6:3044-3066 (1986)) were prepared as described below. Cultures were prepared on five 13 mm glass coverslips, each of which was coated with poly-L-lysine. Five days after plating, cultures were maintained for at least 10 days before infection with pHSVlac. Cultures contained approximately 1×10^5 cells per 35 mm dish at the time of pHSVlac infection.

EXAMPLE 2

CONSTRUCTION AND PROPAGATION OF pHSVlac STOCK

pHSVlac was constructed using standard recombinant DNA techniques (Maniatis, T. et al., Molecular Cloning, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). The SV40 origin region and the 750 bp fragment flanked by Bam HI and Eco RI restriction sites were excised from the plasmid pCH110 (Hall, C.V. et al., J. Molec. and App. Genet 2:101-109

-23-

(1983)). The SV40 origin region was replaced with the HSV-1 c region nucleotides 47 to 1066 (McGeoch, D.J. *et al.*, Nucleic Acids Research 14:1727-1745 (1986)) and the 750 bp fragment flanked by the Bam HI and Eco RI restriction sites was replaced with the a sequence (Davison, A.J. *et al.*, J. Gen. Virol. 55:315-331 (1981)) nucleotides 127 to 1132 to yield the 8.1 kb plasmid pHSVlac.

The vector was packaged into an HSV-1 viral stock in the following manner: 1.5×10^5 CV1 monkey fibroblasts were seeded on a 60 mm plate, and incubated at 37°C. The following day, the cells were transfected (Graham, F.L. *et al.*, Virol. 52:456-467 (1973)) with a 0.5 ml of calcium phosphate coprecipitate containing 1 μ g HSV-1 vector DNA and 9 μ g salmon sperm DNA. Four hours later, the cells were shocked with 15% glycerol (Parker, B.A. *et al.*, J. Virol. 31:360-369 (1979)). Following a 24 hour incubation at 37°C, 1.5×10^6 PFU of HSV-1 in 100 μ l was added to each plate for 1 hour at room temperature. The virus inoculum was aspirated, and the plates were washed once with 5 ml of medium. Semiconfluent plates of CV1 cells which were not exposed to either virus or DNA were trypsinized. 5 ml of medium containing 5×10^6 recently trypsinized CV1 cells was added to complete the monolayer. Following incubation for 1.5 hours at 37°C, to allow the CV1 cells to attach to the plates, the medium was removed and replaced with either fresh liquid medium or methocel. After incubation for 3 days, virus was harvested or after incubation for 5 days, plaques were visualized with neutral red. Virus stocks were subsequently passaged at a 1:2 dilution on CV1 cells. This procedure will aid in propagating many different defective HSV-1 vectors, and in evaluating the properties of different strains and mutants of HSV-1 helper virus.

-24-

EXAMPLE 3
RECOVERY OF pHSVlac FROM NEURONS

5 An experiment was performed to demonstrate that pHSVlac DNA was present in cells two weeks after infection. It has been demonstrated that superinfection of a latently infected neuron results in a lytic infection; both the latent and superinfecting genomes are present in the progeny virus (Lewis, M.E. et al., J. Gen. Virol. 65:215 (1984)). Such an experiment was modified to investigate the question of whether pHSVlac could be recovered from previously infected cells.

10 To accomplish this goal, cultures (Hawrot, E. et al., Methods Enzymol. 58:574 (1979)) of dorsal root ganglia (9 x 10⁴ cells in 1.5 ml) were infected with 0.05 ml pHSVlac virus stock and incubated for two weeks at 37°C. Two weeks after infection with pHSVlac, cultures were infected with strain 17 ts K alone and incubated for two days at 31°C in an effort to recover pHSVlac DNA in a HSV-1 virus stock.

15 Cultures were infected with 5 x 10⁵ pfu of HSV-1 strain 17 ts K (Davison, M.J. et al., J. Gen. Virol., 65:859 (1984)) and incubated for two days at 31°C. The resulting virus stock was passaged three times on 2 x 10⁶ CV1 monkey fibroblasts at 31°C to yield virus stocks DRG1 and DRG2. 1 x 10⁷ CV1 cells were infected with 5 x 10⁷ pfu of virus stock (DRG1, DRG2, ts K alone, or mock infected) and incubated at 31°C for 24 hours.

20 DNA was analyzed as follows: 1 x 10⁷ CV1 cells were infected with 5 x 10⁷ pfu of pHSVlac virus stock and the cells were incubated at 31°C for 24 hours. Total cellular DNA was prepared as described (Wigler, M. et al., Cell 16:777-785 (1979)) from CV1 cells, 5 µg of DNA was digested with 12.5 units of the restriction endonuclease Eco RI overnight at 37°C, resolved on 0.7% agarose gels, and transferred to Genetran as described (Southern, E.M., J. Molec. Biol. 98:503-

25

30

517 (1975)). Hybridization and washing conditions were as described (Southern, E.M. J. Molec. Biol. 98:503-517 (1975)).

5 The probe was the 5.9 kb Eco RI fragment from the plasmid pCH110 (Hall, C.V. et al., J. Molec. and App. Genet 2:101-109 (1983)) radiolabeled with ³²P (Feinberg, A.P. et al., Analytical Biochem. 132:6-13 (1983)). This probe is homologous to pBR322 sequences and most of lacZ, lacking homology to the 133 bp at the 3' end. pHSVlac contains three Eco RI sites, one at each end of the pBR322 segment and a
10 third in the lacZ gene 133 bp from the 3' end of the fragment. The 4.3 kb band contains most of the transcription unit in pHSVlac and the 2.3 kb band contains the pBR322 sequences. These two bands are present in equimolar amounts. A 1.5 kb fragment, which contains the 3' end of the lacZ gene, the SV-
15 40 early region polyadenylation site, and the a sequence is not homologous to the probe. pHSVlac sequences are present in virus stocks containing the vector but are absent from virus stocks of ts K alone and from uninfected cells.

20 The presence of pHSVlac DNA was identified in this manner. The structure of pHSVlac DNA which persisted in sensory neurons for two weeks was found to be similar to the structure of pHSVlac DNA isolated from E. coli. pHSVlac DNA was found to be absent from both a virus stock of strain 17 ts K alone and from mock infected cells. 1 to 10% of PC12 cells
25 were β -galactosidase positive 24 hours after infection with virus stocks of pHSVlac recovered from neurons two weeks after infection. Thus pHSVlac DNA can persist, unaltered, in sensory neurons for at least two weeks and stably express β -galactosidase from the HSV-1 IE 4/5 promoter.

-26-

EXAMPLE 4EXPRESSION OF THE lacZ TRANSCRIPTION UNIT OF pHSVlac IN
MITOTIC CELLS AT THE RESTRICTIVE TEMPERATURE OF 37°C

5 The fate of pHSVlac DNA in cells incubated at the
restriction temperature of 37°C was investigated. CV1 cells
were infected with pHSVlac virus stock, and the cells were
incubated at 37°C. Twenty-four hours after infection total
cellular DNA was isolated and subjected to Southern analysis
as described above. The results demonstrated that pHSVlac DNA
10 is present in CV1 cells incubated for 24 hours at 37°C.
Similar blots probed with an HSV-1 specific probe
demonstrated that HSV-1 DNA was also present in these cells.

15 The ability of pHSVlac to transcribe the lacZ
transcription unit in pHSVlac was investigated. CV1 cells
were infected with pHSVlac virus stock, and the cells were
incubated for 24 hours at 37°C. Total cellular RNA was
prepared (Chirgwin, J.M. et al., Biochemistry 18:5294-5299
(1979)), displayed on agarose gels, and the resulting
Northern blots (Ecker, J.R. et al., Proc. Natl. Acad. Sci. USA
20 84:5202-5206 (1987)) were probed with a lacZ specific probe.
A prominent band seen at 4.0 kb is the appropriate size for
the correctly processed transcript from the lacZ transcription
unit using the HSV-1 1E 4/5 promoter in pHSVlac. This 4 kb
transcript is not present in cells infected with ts K alone
25 nor in uninfected cells; therefore it must be derived from
pHSVlac. Similar blots probed with a HSV-1 specific probe
showed a pattern of bands indicative of expression of the HSV-
1 IE genes in cells infected with pHSVlac and in cells
infected with ts K alone but not in uninfected cells.

-27-

EXAMPLE 5EXPRESSION OF β -GALACTOSIDASE BY pHSVlac

To determine if pHSVlac virus stock could infect neurons and express β -galactosidase, primary cultures of dissociated neurons were prepared (Hawrot, E. et al., Methods Enzymol. 58:574 (1979)), and the expression of β -galactosidase from pHSVlac in cells from dorsal root ganglia and superior cervical ganglia of newborn rats was studied.

The virus stock containing pHSVlac was prepared with HSV-1 strain 17 ts K (Davison, M.J. et al., J. Gen. Virol., 65:859 (1984)) as helper virus. The titer of the virus stock was 1×10^6 pfu of strain 17 virus per ml and 8×10^5 infectious particles of pHSVlac per ml. Cultures were infected with pHSVlac virus stock, then incubated for 24 hours at 37°C, fixed, and assayed for β -galactosidase activity in situ (Sanes, J.R. et al., EMBO J. 5:3133 (1986); Price, J. et al., Proc. Natl. Acad. Sci. USA 84:156 (1987)) using the chromogenic substrate, 5-bromo-4-chloro-3-indoyl β -D-galactoside (X-gal). Dissociated cell cultures (Hawrot, E. et al., Methods Enzymol. 58:574 (1979)) were prepared from newborn rat dorsal root ganglia (DRG) or superior cervical ganglia (SCG) and treated for 24 hours with 10^{-5} M cytosine arabinoside. After 10 days in vitro, the cultures contained 3 to 8×10^5 cells per 35-mm plate; cultures were then infected with 0.1 ml of pHSVlac virus stock and incubated for 24 hours at 37°C. Cells were fixed with 0.5% glutaraldehyde and stained for beta-galactosidase activity with X-gal. The stained preparations were photographed and analyzed.

When infected at a multiplicity of infection (moi) of 0.1 to 0.4 (pHSVlac per cell) about 38% of the cells in the cultures of dorsal root ganglia and 11% of the cells in the cultures of superior cervical ganglia were β -galactosidase positive.

-28-

Experiments performed at higher ratios of pHSVlac virus to cells (moi=2) resulted in expression of β -galactosidase in virtually every cell. Cultures infected with HSV-1 strain 17 ts K alone or mock infected cultures contained less than 0.2% β -galactosidase positive cells.

Most of the β -galactosidase positive cells observed had the morphological characteristics of neurons. However, β -galactosidase positive cells which resembled glia were also observed.

These results demonstrate that the IE 4/5 promoter is capable of mediating constitutive gene expression in infected neurons. The expression of β -galactosidase was found to be independent of the moi of the ts K helper virus. A single particle of pHSVlac was sufficient to render a cell β -galactosidase positive. A productive lytic infection was not required in order to express the β -galactosidase gene.

EXAMPLE 6

NEURONAL INFECTION OF pHSVlac

To determine if some of the β -galactosidase positive cells identified in the experiments of Example 5 were indeed neurons, an experiment was performed to determine if β -galactosidase and a neuronal marker were present in the same cell. To do this, primary cultures of dorsal root ganglia (Hawrot, E. et al., Methods Enzymol. 58:574 (1979)) were infected with pHSVlac virus stock, incubated for 24h at 37°C, fixed, and treated with a rabbit antiserum to β -galactosidase and mouse anti-rat monoclonal to one of two neuronal markers, either the A2B5 antigen (Eisenbarth, G.S. et al., Proc. Natl. Acad. Sci. USA 76:4913 (1979)) or the 150 and 180 kD subunits of neurofilament (Sternberger, L.A. et al., Proc. Natl. Acad. Sci. USA 80:6126 (1983)).

-29-

In detail, pHSVlac virus stock and cultures of dorsal root ganglia were prepared as described above except that cultures were prepared on 13-mm glass coverslips coated with 0.8 μ g of laminin. Cultures (5×10^4 cells in 0.5 ml) were infected with 0.1 ml of pHSVlac virus stock and incubated for 24 h at 37°C. Fixation with 4% paraformaldehyde in 0.1 M NaPO_4 (pH 7.0) and immunohistochemistry was performed (Huettnner, J.E. *et al.*, *J. Neurosci.* 6:3044 (1986)) with a rabbit antibody to *E. coli* β -galactosidase (Cooper Biomedical, Malvern Pa.) diluted 1:800 and either mouse monoclonal to rat neurofilament (Sternberger, L.A. *et al.*, *Proc. Natl. Acad. Sci. USA* 80:6126 (1983)) (SMI 33, Stenberer-Meyer) diluted 1:800 or mouse monoclonal A2B5 (Eisenbarth, G.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 76:4913 (1979)) supernatant diluted 1:2 as primary antibodies. Fluorescein isothiocyanate-conjugated goat F(ab')₂ antibody to mouse F(ab')₂ (Cooper Biomedical) diluted 1:200 and rhodamine isothiocyanate-conjugated goat F(ab')₂ antibody to rabbit F(ab')₂ diluted 1:250 (Cooper Biomedical) were used as secondary antibodies. Coverslips were mounted in PBS glycerol 1:1 containing 0.4% n-propyl gallate. β -galactosidase-like immunoreactivity (β -gal-IR) was visualized with a rhodamine conjugated goat antibody to rabbit immunoglobulin G and A2B5 or neurofilament-like immunoreactivity (A2B5-IR or Nf-IR) was visualized with a fluorescein conjugated goat antibody to mouse immunoglobulin G.

Many of the same cells with neuronal morphology contained both A2B5-IR and β -gal-IR or Nf-IR and β -gal-IR. Parallel cultures treated with antibodies directed against either of the neuronal markers and rabbit preimmune serum followed by the fluorescent-conjugated antibodies contained either A2B5-IR or Nf-IR but no β -gal-IR. Cultures infected with *ts K* alone, or mock infected, and treated with antibodies against either neuronal marker and β -galactosidase contained either A2B5-IR

-30-

or Nf-IR but no β -gal-IR. Thus, pHSVlac can infect rat sensory neurons and subsequently express β -galactosidase.

Example 7

EXPRESSION IN NEURONS

5 Primary cultures were prepared from a variety of CNS
areas, including spinal cord, cerebellum, thalamus, striatum,
10 hippocampus, occipital cortex, temporal cortex, and frontal
cortex. The cultures were grown for at least 10 days and
treated with cytosine arabinoside to kill mitotic cells,
thereby preventing glial cell overgrowth of the cultures.
Cultures were then infected with pHSVlac, and after
incubation at 37°C for either 24 hours or 2 weeks, assays were
performed to detect pHSVlac DNA and expression of E. coli β -
galactosidase in neurons.

15 To demonstrate such expression, cultures were fixed and
assayed for the co-localization of β -galactosidase-like
immunoreactivity and neurofilament-like immunoreactivity using
the immunofluorescent staining assay described above. Phase-
positive cells from occipital cortex, temporal cortex and
20 hippocampus demonstrated staining for both β -galactosidase-
like and neurofilament-like immunoreactivity. In these
cultures, between 60-70% of the neurons in microscopic fields
examined were β -galactosidase positive. Some cells which
contained β -galactosidase-like reactivity lacked
25 neurofilament-like activity, and had the morphological
appearance of glia, although further characterization of these
cells was not made.

30 β -galactosidase-like reactivity was observed in cells
which contained neurofilament-like reactivity in cultures of
spinal cord, cerebellum, thalamus, striatum and frontal
cortex. Sister cultures treated with pre-immune primary sera
contained background levels of fluorescein and rhodamine

-31-

fluorescence, and sister cultures treated with antibody against neurofilament antigens and rabbit pre-immune serum followed by the fluorescent-conjugated antibodies contained neurofilament-like immunoreactivity but did not contain β -galactosidase-like immunoreactivity.

Cultures infected with HSV strain 17 ts K alone, or mock infected, and treated with anti-neurofilament and anti- β -galactosidase antibodies followed by the fluorescent conjugated secondary antibodies, contained neurofilament-like immunoreactivity but did not contain β -galactosidase-like immunoreactivity. Phase-positive cells from the total neocortex and striatum demonstrated both neurofilament-like immunoreactivity and β -galactosidase-like immunoreactivity 2 weeks after infection with pHSVlac. Between 60-70% of striatal and neocortical neurons examined were β -galactosidase positive.

EXAMPLE 8

PERSISTENCE OF pHSVlac AND STABILITY OF EXPRESSION

The persistence of pHSVlac, and the stability of its expression were investigated in cultured sensory neurons. Cultures of dorsal root ganglia (Hawrot, E. et al., Methods Enzymol. 58:574 (1979)) were infected with pHSVlac virus stock; following a two week incubation at 37°C, 49% of the cells were β -galactosidase positive as assayed with X-gal. Most of these cells had neuronal morphology. Cultures infected with HSV-1 strain ts K alone or mock infected contained less than 0.2% β -galactosidase positive cells.

The presence of β -galactosidase positive cells could have resulted from either stable persistence of pHSVlac in the same cell for two weeks or horizontal transmission of pHSVlac. If the presence of beta-galactosidase resulted from the stable

-32-

5 persistence of pHSVlac, then the proportion of total cells expressing beta-galactosidase should have been approximately the same as the proportion of cells initially infected with the pHSVlac vector. In contrast, if horizontal transmission had occurred at a significant rate then all the cells would have contained pHSVlac DNA and would have expressed β -galactosidase, and both 17 ts K virus and pHSVlac virus would have been found in the culture medium.

10 Two weeks after infection, between 30-50% of the cells were found to be β -galactosidase negative, but neurofilament positive by immunofluorescence. Furthermore, two weeks after infection the amount of virus in the culture medium was either below detection (i.e. less than 10 plaque forming units (pfu) of strain 17 ts K per milliliter and less than 10 infectious particles of pHSVlac per milliliter), or very low (Table 1).
15 The low level of virus observed might have been due to the release of virus from dead cells during the freezing and thawing of the preparation. In contrast, wt HSV-1 kills all cells in less than 24 hours.

20 Furthermore, in cultured peripheral nervous system neurons, as well as in cultures of differentiated PC-12 pheochromocytoma and differentiated N1E-115 mouse neuroblastoma cells, pHSVlac infection results in no detectable ts K or pHSVlac virus in the culture medium 2
25 weeks after infection. Thus, although it is not necessarily absent, the rate of horizontal transmission of pHSVlac is very low.

30 The experimental results show that the expression of beta-galactosidase resulted predominately, if not completely, from the persistence of the vector rather than its horizontal transmission.

-33-

TABLE 1
AMOUNT OF ts K AND pHSVlac VIRUS PRESENT IN MEDIUM FROM
PRIMARY CULTURES OF NEOCORTEX TWO-WEEKS POST-INFECTION
WITH pHSVlac

	Culture	ts K /ml ⁺	pHSVlac / ml*
5	1	8×10^2	1×10^1
	2	<10 ¹	4×10^1
	3	<10 ¹	<10 ¹
10	4	<10 ¹	<10 ¹

EXAMPLE 9
DIFFERENTIATED CELL CULTURE

5 x 10⁵ PC12 cells (Greene, L.A. et al., Proc. Natl. Acad. Sci. USA 73:2424-2428 (1976)) were seeded in 5 ml on 60 mm plates coated with 0.2 ml of 100 ug/ml collagen and 12 hours later nerve growth factor (NGF) was added to a final concentration of 10 ng/ml. 5 x 10⁵ N1E-115 cells (Amano, T. et al., Proc. Natl. Acad. Sci. USA 69:258-263 (1972); Garvican, J.H. et al., Eur. J. Biochem. 76:251-261 (1977); Nelson, P. et al., Proc. Natl. Acad. Sci. USA 64:1004-1010 (1969)) were seeded in 5 ml on uncoated 60 mm plates and 12 hours later dibutyryl cyclic AMP (bt₂CAMP) was added to a final concentration of 1 mM. The cultures were fed with media containing the appropriate differentiating agent on day 2. On day 3 cultures were infected with 1 x 10⁵ pfu of pHSVlac virus stock, and incubated for 24 hours or two weeks at 37°C. After the two week incubation, 1 x 10⁶ CV1 cells were added to some N1E-115 cultures which were then incubated for an additional two days at 37°C. Cultures were then assayed for β -galactosidase activity in situ.

Alternatively, cultures were infected with 5 x 10⁵ pfu of HSV-1 strain 17 ts K and incubated for two days at 31°C. The resulting virus stock was passaged three times on 2 x 10⁶ CV1 monkey fibroblasts at 31°C. 1 x 10⁷ CV1 cells were infected

-34-

with 5×10^7 pfu of virus stock and incubated at 31°C for 24 hours, and total cellular DNA was prepared (Wigler, M. et al., Cell 16:777-785 (1979)).

5 The structure of pHSVlac DNA and the helper virus DNA in the virus stock was examined. CV1 cells were infected with pHSVlac virus stock, and the cells were incubated at the permissive temperature of 31°C. Twenty-four hours after infection total cellular DNA was isolated (Wigler, M. et al., Cell 16:777-785 (1979)), and analyzed by Southern analysis as described above.

10 This experiment demonstrated that pHSVlac DNA is faithfully propagated in a HSV-1 virus stock using ts K as helper virus. Similar Southern blots were probed with a HSV-1 and pHSVlac specific probe. These blots demonstrated that the structure of the ts K grown in the presence of pHSVlac is similar to the structure of the ts K grown alone and that HSV-1 sequences are absent from uninfected cells. Furthermore, densitometry scanning of a band specific to pHSVlac and a band specific to ts K demonstrated that pHSVlac is present in roughly equimolar amount to ts K.

EXAMPLE 10

INFECTION OF NON-MITOTIC CELLS WITH PHSVAC RESULTS IN PERSISTENCE OF pHSVAC DNA AND STABLE EXPRESSION OF β -GALACTOSIDASE

25 As described above, pHSVlac can infect rat peripheral neurons and express β -galactosidase in a heterogeneous population of cells. For many types of experiments a homogenous population is required. Two non-mitotic cell culture systems; PC12 rat pheochromocytoma cells
30 differentiated with NGF (Greene, L.A. et al., Proc. Natl. Acad. Sci. USA 73:2424-2428 (1976)) and N1E-115 adrenergic mouse neuroblastoma cells differentiated with bt₂cAMP (Amano,

5 T. et al., Proc. Natl. Acad. Sci. USA 69:258-263 (1972);
Garvican, J.H. et al., Eur. J. Biochem. 76:251-261 (1977);
Nelson, P. et al., Proc. Natl. Acad. Sci. USA 64:1004-1010
(1969)); are homogenous population of cells which closely
10 resemble neurons. wt HSV-1 can infect PC12 cells (Rubenstein,
R. et al., J. Gen. Virol. 64:2505-2509 (1983)) and ts K can
infect and persist in differentiated N1E-115 cells (Gerdes,
J.G. et al., Virol. 94:430-441 (1979)). PC12 and N1E-115
cells were differentiated with the appropriate agent for three
15 days. Cultures were then infected with pHSVlac virus stock,
incubated for 24 hours at 37°C, fixed with 0.5%
glutaraldehyde, and assayed for β -galactosidase activity in
situ, using X-gal (Price, J. et al., Proc. Natl. Acad. Sci.
USA 84:156-160 (1987); Sanes, J.R. et al., EMBO J. 5:3111-3142
20 (1986)); 13% of the PC12 cells and 7% of the N1E-115 cells
were β -galactosidase positive. The long processes in the β -
galactosidase positive cells argue that mitosis had been
arrested and differentiation had occurred before infection
with pHSVlac. Differentiated PC12 cells and N1E-115 cells
25 infected with ts K alone, or uninfected cells, contained less
than 0.2% β -galactosidase positive cells.

To determine if pHSVlac can persist in non-mitotic
cells, differentiated PC12 cells and N1E-115 cells were
infected with pHSVlac virus stock, incubated for two weeks at
25 37°C, and assays were performed to detect pHSVlac DNA and
expression of β -galactosidase. To demonstrate expression of
 β -galactosidase two weeks after infection with pHSVlac,
cultures were fixed, and assayed for β -galactosidase activity
using X-gal, (Price, J. et al., Proc. Natl. Acad. Sci. USA
30 84:156-160 (1987); Sanes, J.R. et al., EMBO J. 5:3111-3142
(1986)). Cultures infected with ts K alone or mock infected
contained less than 0.2% β -galactosidase positive cells

The ability of pHSVlac DNA to persist in non-mitotic PC12
and N1E-115 cells was investigated. It has been established

-36-

that superinfection of a latently infected neuron results in a lytic infection and both the superinfecting genome and the latent genome are present in the progeny virus (Lewis, M.E. et al., J. Gen. Virol. 65:215-219 (1984)). Differentiated PC12 and N1E-115 cells were infected with pHSVlac virus stock and incubated for two weeks at 37°C. Cultures were then infected with ts K alone and incubated for two days at 31°C. Total cellular DNA was isolated (Wigler, M. et al., Cell 16:777-785 (1979)) from the resulting virus stocks, digested with the restriction endonuclease Eco RI, and Southern analysis (Southern, E.M. J. Molec. Biol. 98:503-517 (1975)) was performed to detect pHSVlac DNA as described above. The Southern blot revealed that pHSVlac DNA could faithfully persist in non-mitotic cells two weeks after infection but is absent from virus stocks of ts K alone and from uninfected cells. PC12 cells were infected with the pHSVlac virus stocks recovered from non-mitotic cells two weeks after infection, incubated from 24 hours at 37°C, fixed, and assayed for β -galactosidase activity with X-gal; 1% to 10% of the cells were β -galactosidase positive. In summary, three lines of evidence demonstrate that pHSVlac DNA can persist unaltered in non-mitotic cells for two weeks: First, the recovered pHSVlac DNA has a functional α sequence and HSV-1 origin of DNA replication since it was propagated in a HSV-1 virus stock; second, it has a functional transcription unit which expresses β -galactosidase in both non-mitotic cells two weeks after infection and in PC12 cells following recovery by superinfection; and third, the structure of pHSVlac DNA was unaltered as demonstrated by Southern analysis. Thus, pHSVlac can infect and persist in differentiated PC12 and N1E-115 cells, and stably express β -galactosidase for at least two weeks.

-37-

EXAMPLE 11INFECTION WITH PHSVlac OF A WIDE RANGE OF MITOTIC
CELL TYPES FROM HUMANS, MONKEYS, AND RODENTS RESULTS IN
EXPRESSION B-GALACTOSIDASE

5 The ability of defective HSV-1 vectors to deliver genes
into different mitotic cells types was investigated. A wide
range of immortalized cell lines derived from humans, monkeys,
and rodents were infected with pHSVlac and assayed for
expression of β -galactosidase. The cell lines tested included
10 monkey and mouse fibroblasts, adrenergic and cholinergic mouse
neuroblastoma cells (Amano, T. *et al.*, Proc. Natl. Acad. Sci.
USA 69:258-263 (1972); Garvican, J.H. *et al.*, Eur. J. Biochem.
76:251-261 (1977); Nelson, P. *et al.*, Proc. Natl. Acad. Sci.
USA 64:1004-1010 (1969)), rat pheochromocytoma cells (Greene,
15 L.A. *et al.*, Proc. Natl. Acad. Sci. USA 73:2424-2428 (1976)),
rat and mouse pituicytes (Kitagawa, S. *et al.*, Endocrinology
120:2591-2596 (1987); Herbert, E. *et al.*, Endocrinology
102:218-226 (1978)), and a human neuroblastoma (Biedler, J.L.
et al., Cancer Research 38:3751-3757 (1978)). Cultures were
20 infected with pHSVlac virus stock, incubated for 24 hours at
37°C, and the amount of β -galactosidase activity was
determined by a quantitative solution assay using ONPG.
Parallel cultures were fixed, assayed for β -galactosidase
activity *in situ* using the chromogenic substrate X-gal, and
25 the percentage of β -galactosidase positive cells was
determined. The ratio of β -galactosidase activity in cultures
infected with pHSVlac and ts K was evaluated. This ratio was
normalized to a per cell infected with pHSVlac basis using the
results from the *in situ* assay. As shown in Table 2, high
30 levels of β -galactosidase activity were observed in every cell
line tested. There was a five fold variation in β -
galactosidase levels among the cell lines tested. In CV1
cells 24 hours after infection with pHSVlac the β -

-38-

galactosidase specific activity was 115 nmoles ONPG cleaved/min/mg protein. Thus, it may be concluded that HSV-1 vectors should be useful for transferring genes into a wide range of mitotic mammalian cell lines, including those derived from humans.

TABLE 2
 β -GALACTOSIDASE ACTIVITY PER
 INFECTED CELL IN VARIOUS CELL LINES 24
 HOURS AFTER INFECTION WITH pHSVlac VIRUS

	Cell Line	A ₄₂₀ pHSVlac/ts K
	CV1 Monkey Fibroblast	264
	LM tk ⁻ Mouse Fibroblast	59
	N1E-115 Mouse Adrenergic	
	Neuroblastoma	85
	NS-20Y Mouse Cholinergic	
	Neuroblastoma	65
	PC12 Rat Pheochromocytoma	156
	AtT-20 Mouse Pituicyte	65
	GH4 Rat Pituicyte	78
	SK-N-BE(2) Human Neuroblastoma	52

1 x 10⁶ cells were infected at a multiplicity of 0.1 with pHSVlac virus stock, or with ts K alone. The cells were incubated for 24 hours at 37°C. β -galactosidase activity per cell infected with pHSVlac was determined as described in Experimental Procedures. Each measurement is the average of three separate cultures from at least two separate experiments whose values differed by less than 10%.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth and as follows in the scope of the appended claims.

-39-

International Application No: PCT/

MICROORGANISMS	
Optional Sheet in connection with the microorganism referred to on page <u>4</u> , line <u>29</u> of the description *	
A. IDENTIFICATION OF DEPOSIT *	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution *	
American Type Culture Collection	
Address of depositary institution (including postal code and country) *	
12301 Parklawn Drive Rockville, Maryland 20852, United States of America	
Date of deposit *	Accession Number *
31 January 1989	40544
B. ADDITIONAL INDICATIONS * (leave blank if not applicable). This information is continued on a separate attached sheet <input type="checkbox"/>	
HSV-1 Defective Virus Plasmid DNA, pHSVlac	
C. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE * (if the indications are not for all designated States)	
D. SEPARATE FURNISHING OF INDICATIONS * (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later * (Specify the general nature of the indications e.g. "Accession Number of Deposit")	
E. <input checked="" type="checkbox"/> This sheet was received with the international application when filed (to be checked by the receiving Office)	
(Authorized Officer) <i>Theodore L. Sigler</i>	
<input type="checkbox"/> The date of receipt (from the applicant) by the International Bureau **	
(Authorized Officer)	

(January 1985)

-40-

WHAT IS CLAIMED IS:


1. A recombinant HSV-1 vector capable of infecting a neuronal cell, wherein said vector contains:
 - (1) an HSV-1 packaging site-containing sequence;
 - 5 (2) an HSV-1 origin of DNA replication-containing sequence; and
 - (3) a heterologous reporter or selectable marker gene sequence sufficient to permit the recognition or selection of the vector.
- 10 2. The recombinant vector of claim 1 which additionally contains:
 - (4) a promoter sequence sufficient to direct transcription of a distally located sequence in a neuronal cell;
 - 15 (5) a sequence sufficient to direct translation of a distally located sequence in a neuronal cell; and
 - (6) a sequence sufficient to permit a neuronal cell to process an expressed protein.
- 20 3. The recombinant vector of claim 2 which additionally contains:
 - (7) a sequence sufficient to permit the propagation of the vector in a bacteria; and
- 25 4. The recombinant vector of claim 3 which additionally contains:
 - (8) an inserted gene sequence;
wherein said inserted gene sequence is inserted into said vector such that it is operably linked to said sequences (4), (5) and (6).

-41-

5. The recombinant molecule of any of claims 1 or 4 wherein said HSV-1 packaging site-containing sequence (1) contains the HSV-1 a segment.
- 5 6. The recombinant molecule of any of claims 1 or 4 wherein said HSV-1 origin of DNA replication sequence (2) contains the HSV-1 ori_S region.
7. The recombinant molecule of any of claims 1 or 4 wherein said heterologous reporter or selectable marker gene sequence (3) contains a sequence from the E. coli β -galactosidase gene.
- 10 8. The recombinant molecule of any of claims 2 or 4 wherein said promoter sequence (4) contains the HSV-1 IE 4/5 promoter sequence.
9. The recombinant molecule of any of claims 2 or 4 wherein said
- 15 sequence (5) sufficient to direct translation of a distally located sequence in a neuronal cell contains the HSV-1 intron sequence which follows the HSV-1 IE 4/5 promoter sequence.
10. The recombinant molecule of any of claims 2 or 4 wherein said
- 20 sequence (6) sufficient to permit a neuronal cell to process an expressed protein contains the SV40 early region polyadenylation site.
11. The recombinant molecule of claim 1 which is capable of persisting in a non-mitotic cell.
- 25 12. The recombinant molecule of claim 1 which is pHSVlac.

-42-

13. The recombinant molecule of claim 4 wherein said inserted gene sequence (8) is a gene sequence associated with a disease selected from the group consisting of: lysosomal storage disease, the Lesch-Nyhan syndrome, amyloid polyneuropathy, Alzheimer amyloid, and Duchenne's muscular dystrophy.



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US90/00357

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *
According to International Patent Classification (IPC) or to both National Classification and IPC
IPC(5): C 12 N 15/00, 7/00; C 12 P 21/00
U.S. 435/172.3, 172.1, 235, 69.1

II. FIELDS SEARCHED

Classification System	Minimum Documentation Searched ?
U S	435/172.3, 172.1, 235, 69.1,

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched *

Cas database (1967-1990) keywords: herpes simplex virus/
vector/recombinant /express ?/ gene(s)

III. DOCUMENTS CONSIDERED TO BE RELEVANT *

Category *	Citation of Document, * with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **
X Y	SHIH, M. ET AL. (SEPTEMBER 1984) Proc. Natl. Acad. Sci. USA. Vol. 81: 5867-5870. "Expression of HEPATITIS B VIRUS S GENE by HERPES SIMPLEX VIRUS type 1 vectors carrying alpha and beta regulated gene chimeras." See entire document.	1,2,4-6,11 3,7-10,12,13
X Y	PALLELLA, T. ET AL. (JANUARY 1988) Mol. Cell Biol. Vol 8, No. 2 457-460 "Herpes Simplex Virus-Mediated HUMAN HYPOXANTHINE-GUANINE Phosphoribosyltransferase Gene Transfer Into Normal Cells". See entire document.	1,2,4-6,11,13 3,7-10,12
X P	Chemical Abstracts, Volume 112 No. 5 issued 29 JANUARY 1990 (COLUMBUS, OHIO, USA) GELLER, A ET AL. "Herpes Simplex virus 1 (HSV1) vector system for introduction of foreign genes to rat brain neuron." See page 128, column 1, abstract number 312419 JIKKEN IGAKU 1989, 7(16), 1941-53.	1-13
P,X	PALLELA, T. D. ET AL. (AUGUST 1989) Gene, volume 80: 137-144 "Expression of human HPRT MRNA in brains of mice infected with a recombinant herpes simplex virus-1 vector." See entire document.	1,2,4-6,11,13
(See attachments)		

* Special categories of cited documents: *

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

30 APRIL 1990

International Searching Authority

ISA/US

Date of Mailing of this International Search Report

17 JUL 1990

Signature of Authorized Officer

ANNE BROWN

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
P,X	Chemical Abstracts, Volume 110, NO. 7 issued 13 FEBRUARY 1989 (Columbus, Ohio, USA) MEIGNIER B. ET AL. " in vivo behavior of genetically engineered herpes simplex viruses R7017 and R7020: construction and evaluation in rodents." See page 156, column 2 and page 157, column 1. Abstract number 520215, J. Infect. Dis. 1988,158(3) 602-14.	1,2,4,6,11
P,X	Chemical Abstracts, Volume 111, No. 9 issued 28 AUGUST 1989 (Columbus, Ohio, USA) MEIGNIER, B. ET AL. " Genetic Engineering Of Novel Herpes Simplex Virus (HSV) Genomes For Use As Potential Vaccines And As Vectors Of Foreign Genes". See page 187, column 2, Abstract Number 71936F, Int. Biotechnol. Symp. 8th 1988, 2, 740-5.	1,2,4-6,11
X Y	SMILEY, J. ET AL. (AUGUST 1987) J. VIROL. Volume 61: 2368-2377. " Expression of a Cellular Gene Cloned in Herpes Simplex Virus: Rabbit beta globin is regulated as an early viral gene in infected fibroblasts." See entire document.	1,2,4-6,11 3,7-10,12,13